

INSTRUCTIONS FOR USE

ZIMHI™ (ZIM-hye)

(naloxone hydrochloride injection) Prefilled syringe

Use ZIMHI for known or suspected opioid overdose in adults and children.

ZIMHI is intended to be administered by people 12 years of age or older. Younger children or those with limited hand strength may find the device difficult to use.

Read this Instructions for Use carefully before you use this product.

Before you need to use your ZIMHI syringe make sure your healthcare provider shows you the right way to use it. Parents, caregivers, and others who may be in a position to administer ZIMHI should also understand how to use it. If you have questions, ask your healthcare provider.

Get Ready to Use ZIMHI™

ZIMHI works like a standard prefilled syringe.



ZIMHI is injected downwards, into the middle of the outer thigh, (as shown), through clothing if needed.



Ready to Use ZIMHI



Place the patient on their back. When ready to inject, pull off cap to expose needle.



Do not put finger on top of the device.

In children under the age of 1 year old, pinch the thigh muscle while administering the dose. Hold ZIMHI by finger grips only and slowly insert the needle into the thigh.



After needle is in thigh: Push the plunger all the way down until it **clicks** and hold for 2 seconds.



After Use

Right after the injection, using one hand with fingers behind the needle, slide the safety guard over the needle. Do not use two hands to activate the safety guard. Put the used syringe into the blue case and close the case.



Get Help
Get emergency medical help now




9-1-1

Tell healthcare provider that you gave an injection of naloxone hydrochloride. Turn the patient on their side (recovery position) after giving ZIMHI.




Use second syringe if necessary



You may need a second ZIMHI syringe if symptoms continue or return.

If symptoms return after an injection with ZIMHI, an additional injection using another ZIMHI prefilled syringe may be needed. Give additional injections using a new ZIMHI prefilled syringe every 2 to 3 minutes and continue to closely watch the person until emergency help is received. Using ZIMHI does not take the place of emergency medical care.



After Use and Disposal

- ZIMHI is a single-dose prefilled syringe that delivers a fixed dose of naloxone hydrochloride. The prefilled syringe cannot be reused. It is normal for most of the medicine to remain in the syringe after the dose has been injected.
- The correct dose has been injected if the plunger has been pushed all the way down and the solution window is at least partially blocked.
- Tell the healthcare provider that you have received or administered an injection of naloxone hydrochloride. Show the healthcare provider where the injection was administered.
- Give your used ZIMHI syringe, contained in the blue case, to the healthcare provider for inspection and proper disposal.

For more information see Patient Information sheet or ask your healthcare provider.

How to Store — Keep ZIMHI in its plastic case nearby and ready for use at all times.

- Store at controlled room temperature, between 68°F to 77°F (20°C to 25°C). **Do not** expose to extreme cold or heat. For example, **do not** store in your vehicle's glove box and **do not** store in the refrigerator or freezer.
- Store the ZIMHI syringe in its closed plastic case to protect from light.

Check Your Stored ZIMHI Often - The solution should be clear when viewed through the window on the device. If the solution is discolored (yellowish or brown color), cloudy or contains particles, replace the ZIMHI with a new one.

Your ZIMHI Has an Expiration Date — Example YYYY-MM. Replace ZIMHI before the last day of expiration month. Dispose of expired ZIMHI properly by taking the ZIMHI™ in its case to a healthcare provider or hospital emergency room.

For more information about ZIMHI prefilled syringe and proper use of the ZIMHI, call 1-800-230-3935 or visit www.ZIMHI.com.

ZIMHI is a registered trademark of Adamis Pharmaceuticals Corporation.

Manufactured for Adamis Pharmaceuticals, San Diego, CA 92130, USA.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Issued: Month Year



ZIMHI™

(naloxone hydrochloride injection)

Prefilled syringe

INSTRUCTIONS FOR USE

Pull out and Read before Use

Rx Only